

1. Ecology of Peatlands:

Baird, A. J., P. A. Eades, and B. W. J. Surridge .2008. The hydraulic structure of a raised bog and its implications for ecohydrological modelling of bog development, *Ecohydrology*, 1: 289–298.

Hooijer, A., S. Page , J. G. Canadell , M. Silvius , J. Kwadijk , H. Wosten, and J. Jauhiainen. 2010. Current and future CO₂ emissions from drained peatlands in Southeast Asia. *Biogeosciences*, 7, 1505–1514 doi:10.5194/bg-7-1505-2010

Morris, P. J., J. M. Waddington, B. W. Benscoter, and M. R. Turetsky. 2011. Conceptual frameworks in peatland ecohydrology: Looking beyond the two-layered (acrotelm–catotelm) model, *Ecohydrology*, 4(1), 1–11.

Wüst, R. A. J., R. M. Bustin, and L. M. Lavkulich. 2003. New classification systems for tropical organic-rich deposits based on studies of the Tasek Bera Basin, Malaysia, *Catena*, 53(2), 133–163.

Yu, Z. C. 2012. Northern peatland carbon stocks and dynamics: A review, *Biogeosciences*, 9: 4071–4085

Zhao, M. and S.W. Running. 2010. Drought-induced reduction in global terrestrial net primary production from 2000 through 2009. *Science* 329:940 – 943.

2. Measuring & Mapping Depth of Peatlands and carbon stocks using GPR:

Comas, X., et al. 2015. Imaging tropical peatlands in Indonesia using ground-penetrating radar (GPR) and electrical resistivity imaging (ERI): implications for carbon stock estimates and peat soil characterization. *Biogeosciences*, 12: 2995–3007

Hengl,T. et al. 2014. SoilGrids1km—Global Soil Information Based on Automated Mapping. *PLoS ONE*. 2014 08; 9(e105992). doi: [10.1371/journal.pone.0105992](https://doi.org/10.1371/journal.pone.0105992) PMID: [25171179](https://pubmed.ncbi.nlm.nih.gov/25171179/)

Jaenicke, J. et al. 2008. Determination of the amount of carbon stored in Indonesian peatlands. *Geoderma* 147: 151-158.

Kettridge, N., X. Comas, A. Baird, L. Slater, M. Strack, D. Thompson, H. Jol, and A. Binley. 2008. Ecohydrologically important subsurface structures in peatlands revealed by ground-penetrating radar and complex conductivity surveys *Journal of Geophysical Research* 113, G04030, doi:10.1029/2008JG000787.

Kettridge, N., A. Binley, X. Comas, N. J. Cassidy, A. J. Baird, A. Harris, J. van der Kruk, M. Strack, A. M. Milner, and J. M. Waddington. 2012. Do peatland microforms move through time? Examining the developmental history of a patterned peatland using ground-penetrating radar, *Journal of Geophysical Research* 117, G03030, doi:10.1029/2011JG001876.

Loisel, J., Z. Yu, A. Parsekian, J. Nolan, and L. Slater. 2013. Quantifying landscape morphology influence on peatland lateral expansion using ground penetrating radar (GPR) and peat core analysis, *J. Geophys. Res. Biogeosci.*, 118, 373–384, doi:10.1002/jgrg.20029.

Lowry, C. S., D. Fratta, and M. P. Anderson. 2009. Ground penetrating radar and spring formation in a groundwater dominated peat wetland, *J. Hydrol.*, 373(1–2), 68–79.

Parry, L. E., and D. J. Charman. 2013. Modelling soil organic carbon distribution in blanket peatlands at a landscape scale, *Geoderma*, 211–212, 75–84.

Parry, L. E., D. J. Charman, and J. P. W. Noades. 2012. A method for modelling peat depth in blanket peatlands, *Soil Use Manage.*, 28(4), 614–624.

Parry, L.E., L.J. West, J. Holden, and P.J. Chapman. 2014. Evaluating approaches for estimating peat depth. *Journal of Geophysical Research: Biogeosciences* 119: 567-576.

Robinson, D. A., et al. 2008. Soil Moisture Measurement for Ecological and Hydrological Watershed-Scale Observatories: A Review. *Vadoze Zone Journal* 7:358–389

Rosa, E., M. Larocque, S. Pellerin, S. Gagne, and R. Fournier. 2009. Determining the number of manual measurements required to improve peat thickness estimations by ground penetrating radar, *Earth Surf. Processes Landforms*, 34377–383.

Rudiyanto, B. et al. 2016. Digital mapping for cost-effective and accurate prediction of the depth and carbon stocks in Indonesian peatlands. *Geoderma* 2712:20-31.

Sorensen, K.W. 1993. Indonesian peat swamp forests and their role as a carbon sink. *Chemosphere* 27:1065-1082.

Theimer, B. D., D. C. Nobes, and B. G. Warner. 1994. A study of the geoelectrical properties of peatlands and their influence on ground penetrating radar surveying. *Geophysical Prospect.* 42: 179–209.

3. Mapping of Forest Carbon using Active Sensors:

Baccini A, Goetz SJ, Walker WS, Laporte NT, Sun M, Sulla-Menashe D, Hackler J, Beck PSA, Dubayah R, Friedl MA, et al. 2012. Estimated carbon dioxide emissions from tropical deforestation improved by carbon-density maps. *Nature Clim Change*, 2:182–185.

Fatoyinbo, T. E., M. Simard, R. A. Washington-Allen, and H. H. Shugart. 2008. Landscape-scale extent, height, biomass, and carbon estimation of Mozambique's mangrove forests with Landsat ETM+ and Shuttle Radar Topography Mission elevation data, *Journal of Geophysical Research* 113:G02S06, doi:10.1029/2007JG000551.

Fatoyinbo, T. E. & M. Simard .2013. Height and biomass of mangroves in Africa from ICESat/GLAS and SRTM, *International Journal of Remote Sensing*, 34:2, 668-681, DOI:[10.1080/01431161.2012.712224](https://doi.org/10.1080/01431161.2012.712224)

Mitchard et al. 2013. Uncertainty in the spatial distribution of tropical forest biomass: a comparison of pan-tropical maps. **Carbon Balance & Management** 8:10 <http://www.cbmjournals.com/content/8/1/10>

Saatchi et al. 2011. Benchmark map of forest carbon stocks in tropical regions across three continents. *PNAS* 108:9899-9904

4. Mapping Forest Change in Indonesia using passive optical remote sensing:

Hansen, M.C. et al. 2013. High-Resolution Global Maps of 21st-Century Forest Cover Change. *Science* 342:850 DOI: 10.1126/science.1244693

Hansen, M.C., et al. 2009. Quantifying changes in rates of forest clearing in Indonesia from 1990 to 2005 using remotely sensed data sets. *Environ. Res. Lett.*, 4 (034001)

Lepers, E., E. F. Lambin, A. C. Janetos, R. DeFries, F. Achard, N. Ramankutty, and R. J. Scholes. 2005. A synthesis of information on rapid land-cover change for the period 1981–2000. *BioScience* 55:115-124.

5. Mapping of Forest Carbon using Passive Sensors:

Nemani, R., C. Keeling, H. Hashimoto, W. Jolly, , S. Piper, C. Tucker, R. Myneni, S. Running. 2003. Climate-driven increases in global terrestrial net primary production from 1982 to 1999. *Science* 300,1,560-1,563

Nocita M, et al. 2014. Soil spectroscopy: an opportunity to be seized. *Global Change Biology* 21:10–11. doi: [10.1111/gcb.12632](https://doi.org/10.1111/gcb.12632) PMID:24832363

Running, S.W. 2012. A measurable planetary boundary for the biosphere. *Science* 337(6101):1458-1459. DOI: 10.1126/science.1227620

Running, S.W., R.R. Nemani, F.A. Heinsch, M. Zhao, M. Reeves, and H. Hashimoto. 2004. A continuous satellite-derived measure of global terrestrial primary production. *BioScience* 54:547-560.

Sexton, J.O. et al. 2013. Global, 30-m resolution continuous fields of tree cover: Landsat-based rescaling of MODIS vegetation continuous fields with lidar-based estimates of error. *International Journal of Digital Earth* 6(5):427–448. doi: [10.1080/17538947.2013.786146](https://doi.org/10.1080/17538947.2013.786146)

6. GIS & Remote Sensing Mapping of Peatlands:

Connolly, J., N.M. Holden, and S.M. Ward. 2007. Mapping Peatlands in Ireland using a Rule-Based Methodology and Digital Data. *Soil Sci. Soc. Am. J.* 71:492–499 doi:10.2136/sssaj2006.0033

Gou, S.i., S. Gonzales, & G.R. Miller. 2015. Mapping potential groundwater-dependent ecosystems for sustainable management. *Groundwater* 53:99-110.

Jaenicke, J. et al. 2011. Monitoring the effect of restoration measures in Indonesian peatlands by radar satellite imagery. *Journal of Environmental Management* 92:630-638.

Harris, et al. 2015. Hyperspectral remote sensing of peatland floristic gradients. *Remote Sensing of Environment* 162:99 – 111

Krankina, O.N., D. Pflugmacher, M. Friedl, W. B. Cohen, P. Nelson , and A. Baccini. 2008. Meeting the challenge of mapping peatlands with remotely sensed data. *Biogeosciences*, 5:1809–1820

Pflugmacher, D, O.N. Krankina, and W.B. Cohen.2007. Satellite-based peatland mapping: Potential of the MODIS sensor. *Global and Planetary Change* 56: 248–257

Worsfold, R. D., S. K. Parashar, and T. Perrott .1986. Depth profiling of peat deposits with impulse radar, *Can. Geotech. J.*, 23(2), 142–154.

7. RADAR Studies for Biomass:

Imhoff, M.L. et al. 2000. An airborne VHF multi-band SAR for vegetation biomass measurement. *IEEE Trans. Geoscience and Remote Sens.* 38:1458-1462.

Tighe, M.L., D. Chamberlain, G. Lawrence. 2010. What does it take to map the continental U.S. at a high accuracy? *Photogrammetric Engineering and Remote Sensing* 76: 641-643.

8. Websites:

Green Stimulus peat depth survey: <http://www.snh.gov.uk/docs/A1142813.pdf>

Producing a peat volume map for England:

<http://www.northpennines.org.uk/Lists/DocumentLibrary/Attachments/240//Poster.pdf>

International Peatland Society: <http://www.peatsociety.org/peatlands-and-peat/peatlands-and-climate-change>

Peat and Peatlands: <http://www.eolss.net/sample-chapters/c08/e3-04-06.pdf>

1990 Global peat distribution: http://www.grida.no/graphicslib/detail/peat-distribution-in-the-world_8660